

BEGINNER PLUS

22-20 AM DX MOS

Calf Milk Replacer

Type C Medicated Feed

Complete Milk Replacer to be fed to herd replacement calves.

For the prevention of coccidiosis in ruminating and non-ruminating calves (including veal calves) and cattle caused by Eimeria bovis and Eimeria zuernii. Feed as directed.

ACTIVE DRUG INGREDIENT(S)

Decoquinatate 45.4 g/ton
(22.70 mg/lb)

GUARANTEED ANALYSIS

Crude Protein	, Min	22.00 %
Crude Fat	, Min	20.00 %
Crude Fiber	, Max	0.15 %
Calcium	, Min	0.75 %
Calcium	, Max	1.25 %
Phosphorus	, Min	0.70 %
Vitamin A	, Min	30000 IU/lb.
Vitamin D3	, Min	10000 IU/lb.
Vitamin E	, Min	100 IU/lb.

INGREDIENTS

Dried whey product, dried whey protein concentrate, animal fat (preserved with BHA & BHT), dried whey, dried skimmed milk, lecithin, L-lysine, calcium carbonate, brewer's dried yeast, hydrolyzed yeast, DL-methionine, calcium silicate, choline chloride, vitamin E supplement, ferrous sulfate, natural and artificial flavor, zinc sulfate, vitamin A supplement, manganese sulfate, ethoxylated mono and diglycerides, propylene glycol, copper sulfate, ascorbic acid, vitamin B12 supplement, niacin supplement, biotin, vitamin D3 supplement, selenium yeast, calcium pantothenate, riboflavin supplement, menadione sodium bisulfite complex (source of vitamin K activity), pyridoxine hydrochloride, thiamine mononitrate, folic acid, ethylenediamine dihydriodide, cobalt sulfate

Important: Follow mixing and feeding directions.

* * * W A R N I N G : * * *

Do not feed to cows producing milk for food.

C A U T I O N

Do not use in feeds containing bentonite.

Manufactured for:
HUBBARD FEEDS
Mankato, MN
www.hubbardfeeds.com

NET WEIGHT 50 LB (22.67 kg)

01-9986

** MIXING INSTRUCTIONS **

Measure 1.5 quarts of clean, warm (110°F) water into a clean bucket. Fill enclosed cup* with milk replacer powder. Sprinkle 10 oz. of milk replacer powder onto the water surface, mix thoroughly, then add clean, warm water (110°F) until final solution volume reaches 2 quarts. Feed this amount TWICE a day.

*Enclosed cup holds approximately 10 ounces of dry milk replacer powder, weighing powder improves consistency.

** FEEDING INSTRUCTIONS **

Feed at a rate of 22.7 mg decoquinatate per 100 lb of body weight (0.5 mg/kg) per day. Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard.

Based on an 125# calf. Adjust amount fed based on calf body weight.

BIRTH through DAY 1:

Feed 4 quarts of clean, disease-free, high quality maternal colostrum or 1 package of OptiPrime calf colostrum replacer within 4 hours of birth. Feed an additional 2 quarts of colostrum or begin feeding milk replacer within 12 hours.

DAY 2 to WEANING:

Feed 2 quarts of milk replacer solution (100°F) to calves twice a day. Feed 2 quarts once daily for 1 week prior to weaning. Offer free choice clean water and starter daily.

COLD WEATHER FEEDING:

Calves can experience cold stress at temperatures below 50°F. Feed up to 50% more of a NON-MEDICATED milk replacer solution per calf per day or supplement with Calf-Cal high energy supplement when cold stress is a concern.

** MANAGEMENT TIPS **

- Sprinkle milk replacer powder onto water for best mixing.
- Agitate powder and water with a wire whisk.
- Clean and sanitize feeding and mixing equipment after each feeding.
- Introduce a high quality calf starter at 4 days of age.
- Offer clean, fresh water at all times.
- Avoid feeding hay prior to weaning.
- WEAN CALVES AT 5-6 WEEKS OF AGE WHEN EATING A MINIMUM OF 3 LBS. PER HEAD PER DAY OF CALF STARTER FOR 3 CONSECUTIVE DAYS.

Hubbard does not use "Restricted-use Proteins" in their products and is in compliance with FDA and state requirements regarding the use, handling, and storage of "Restricted-use Protein" products.

THIS PRODUCT IS FORMULATED SPECIFICALLY FOR CALVES AND SHOULD BE FED ONLY TO CALVES!

Visit us online at WWW.HUBBARDFEEDS.COM.



51577

REV. 0617